

Granulating Wounds - Instructions for Wounds Healing Naturally

WHAT DO YOU DO FOR...

WOUND CARE:

- **Leave the pressure dressing ON and DRY for AT LEAST 48 HOURS.**
 - Pressure dressings are intentionally tight to decrease bleeding, swelling, and pain.
- After 2 days, to remove the pressure dressing, it is often easiest to soak it in the shower/bath to loosen the adhesive. Once removed, gently clean the area with regular soap and water. You can use baby shampoo when washing your hair for wounds on the scalp.
- **DO NOT** scrub the area, and **DO NOT** allow the shower pressure to hit the area directly.
- Apply **ONLY PLAIN PETROLATUM** to the surgical site **at least once daily** for at least 3 weeks or until fully healed. Try to avoid a scab from forming.
 - Wounds that are kept moist heal faster and leave a better scar than wounds that scab over.
 - Do NOT use any antibacterial ointment because they can cause allergic reactions (unless instructed by your physician).
- It is best to discuss whether to re-bandage the area with your doctor.
 - It may be okay to keep the wound open to air while at home. Please cover the wound when you leave your home as it could be exposed to dust or dirt (e.g. while outdoors).
 - If you prefer to keep the wound covered, you can cover the plain petrolatum with non-adhesive or non-sick gauze and secure it with paper tape. If on your legs, you can avoid using any tape or adhesive, and instead use a sock, stocking, or elastic wrap (e.g., Coban™) to keep the gauze in place.
- **Always practice good hand hygiene** - wash your hands thoroughly with soap and water before changing the dressing or touching your wound to apply plain petrolatum. You can also consider using disposable gloves while performing wound care.

PAIN:

- Apply **ice packs** for at least 5-10 minutes out of every waking hour for 1-2 days after surgery directly over the bandage to reduce swelling and pain. You do not need to apply ice packs while sleeping.
- **Elevate** the surgical site as much as possible to minimize swelling.
 - If you had surgery on your head or neck, relaxing in a recliner or sleeping with extra pillows to prop you up may be helpful.
 - If you had surgery on an arm or hand, you can consider using a sling to help.
 - If you had surgery on a leg, in addition to elevation, your doctor may recommend *compression stockings* when you are on your feet.
- For pain/discomfort, we recommend *acetaminophen* or *ibuprofen*.
 - You may take up to 2 extra strength acetaminophen (1000 mg) and repeat every 8 hours. If you need pain relief in between, you can take 2 ibuprofen tablets (400 mg) and repeat every 6 hours.
 - If your doctor has given you a prescription pain medication, it may already contain acetaminophen and additional acetaminophen should not be taken.
 - **Do not exceed 4000mg of acetaminophen or 3200mg of ibuprofen in a 24-hour period!**
- For any non-resolving pain, contact your doctor.

OPTIMAL HEALING:

- Avoid ANY strenuous activity (i.e. heavy lifting, bending over, or exercise) for at least two days to minimize bleeding risk.
 - Strenuous activity includes running, weightlifting, biking, yoga, elliptical, rowing, & stretching.
- Avoid smoking for at least 3 weeks as it leads to poor wound healing. It is best to stop smoking overall.

WHAT SHOULD YOU EXPECT...

INITIALLY:

- **Bruising, swelling and some pain** are expected after surgery. These will typically resolve in 1-2 weeks. Wounds on the hands, legs and feet may take even longer to improve.
- The more active you are, the more the wound is likely to swell and cause pain.
- Swelling and discomfort are especially common for healing wounds on the hands and legs as swelling tends to pool in these areas because of gravity.
- Your wound may feel **tight, itchy or numb**. This should gradually improve over several months.
- You may notice small amounts of clear or golden-tinged drainage from the wound until the wound is fully healed. This drainage is normal from the healing tissue and from any residual swelling.
- Wounds healing naturally may take weeks to months to fully heal.
 - Wounds on the scalp and hands may require plain petrolatum for 4-6 weeks or longer.
 - Wounds on the legs may require plain petrolatum for 8-12 weeks or longer.

LONG-TERM:

In terms of your final scar, everyone is different and follows a different time course of wound healing. It may take up to **12-18 months to see what the final scar** will look like.

- ONLY silicone-based scar products have proven data to support using them (if desired). We do not endorse one brand over another, but examples include ScarAway®, ScarGuard®, and Kelo-cote®. These types of products are easily found in most pharmacies or online. Do not start applying them until at least 2 weeks after your surgery, or unless otherwise directed by your provider. DO NOT use other products like Mederma® or Vitamin E gels.

If you have any questions/concerns, discuss with your doctor. He/she may make some recommendations to help with the final scar appearance.

WHAT SHOULD YOU DO IF YOU EXPERIENCE...

BLEEDING:

- The pressure dressing over your wound helps to stop bleeding. Any bleeding that you notice can usually easily be stopped with direct firm pressure. DO NOT remove the dressing, elevate the site, and apply **constant pressure** over the dressing for 20 minutes without checking.
- If the pressure dressing becomes completely saturated with blood or if you experience active bleeding that does not resolve with 20 minutes of pressure, **call your doctor's office immediately**.

CONCERNS FOR INFECTION:

- If you experience signs of infection such as increased redness, swelling, warmth, yellow drainage, or worsening pain to touch, call your doctor's office immediately.
 - Some redness over and along the suture line is normal and expected.

WHAT SHOULD YOU DO FOR LONG-TERM SKIN HEALTH:

- Wear sunscreen (broad spectrum for UVA & UVB coverage and SPF≥30), a wide-brimmed hat, and sun protective clothing to avoid future skin cancers. These also help the redness from the surgical scar fade faster.
- See your dermatologist or physician regularly for a complete skin check. Regular skin checks are important for early detection and prevention of skin cancer.